



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

The red powder was boiled for some time with a potassium hydroxide solution, the latter concentrated and acidified with hydrochloric acid, when a yellow mass was precipitated. This was dissolved in hot water, from which it crystallized, on cooling, in yellow leaflets. These melted constantly at  $230^{\circ}\text{C}$ . By reference to *Berichte*, 20, 404, it will be seen that four distinct, nitro-m-oxybenzoic acids exist; of these the  $\beta$ -acid melts at  $230^{\circ}\text{C}$ ., and crystallizes in yellow leaflets. Other points of similarity show that the acid obtained by me is identical with that termed  $\beta$  nitro m-oxybenzoic acid.

The red oil exhibited no signs of crystallization, although I allowed it to stand undisturbed for several months. It was also boiled with concentrated potassium hydroxide for ten hours. After concentration it was allowed to cool, when rather large and well-defined monoclinic prisms, having a deep chrome-red color, appeared. The salt was purified by recrystallization from water, and analyzed. Two estimations of the potassium gave 13.02% and 13.00% K. On heating the salt explodes with violence. The free acid crystallizes from water in long needles having a light yellow color. It melts at  $111^{\circ}\text{C}$ . Like its salt it explodes when heated. Its taste is intensely bitter. In all respects it resembles the trinitro-m-oxybenzoic acid mentioned by Griess, *Annalen*, 117, 28, and Beilstein, *Annalen*, 139, 11. These chemists, however, make no mention of the melting point. The percentage of potassium required by an anhydrous salt of this acid is 12.58%.

The difficulty experienced in obtaining large quantities of the acid is due to the fact that there are other products formed in the nitration process, and these cling tenaciously to the acid, defying the most persistent efforts to effect their removal.

---

*Stated Meeting, May 18, 1888.*

Present, 15 members.

President, Mr. FRALEY, in the Chair.

Correspondence was submitted as follows:

Letters of envoy from Institut Egyptien, Cairo; Institut Météorologique de Roumanie, Bucharest; Museum of Comparative Zoölogy, Cambridge, Mass.

Letters of acknowledgment from Institut Egyptien, Cairo (125); Magyar Tudományos Akademia, Buda-Pesth (125); Prof. Edward Suess, Vienna (125, 126); Verein für Geogra-

phie und Statistik, Frankfurt-am-Main (126); Prof. Paul Albrecht, Hamburg (126); Société Entomologique, Bruxelles (126); Prof. W. B. Dawkins, Manchester, Eng. (126); Mr. Joseph Prestwich, Shoreham, Kent, Eng. (125, 126).

Accessions to the Library were reported from Institut Egyptien, Cairo; Institut Météorologique de Roumanie, Bucharest; K. P. Akademie der Wissenschaften, Berlin; K. Gesellschaft der Wissenschaften, Göttingen; K. Sächsische Gesellschaft der Wissenschaften, Leipzig; R. Accademia dei Lincei, Biblioteca N. C. V. E., Rome; R. Geological Society of Cornwall, Penzance; Mr. R. A. Macfea, Edinburgh; Messrs. J. R. Leeson & Co., Boston; New York Academy of Sciences, Dr. J. S. Newberry, Publishers of "The Globe," New York; Mr. W. J. Potts, Camden; College of Pharmacy, Zoölogical Society, Dr. J. Cheston Morris, Philadelphia; Pennsylvania Geological Survey, Harrisburg; Departments of State and of the Interior, Washington, D. C.; Mr. M. C. Read, Hudson, O.; Museo Michoacano, Morelia, Mexico.

The death of Dr. Gerhard vom Rath was announced as having occurred on April 23, 1888.

Pending nominations Nos. 1176 to 1179 were read and spoken to and balloted for.

Pending nomination No. 1180 was postponed.

The Proceedings of Officers and Council were submitted.

The following persons were reported as having been duly elected members of the Society:

Mr. Talcott Williams, Philadelphia,

Prof. J. McKeen Cattell, Philadelphia,

Prof. Alphonse Favre, Geneva.

And the Society was adjourned by the President.